

Zero temperature transition from d-wave superconductor to underdoped regime

Jinwu Ye ^{1,2} and A. Millis ^{1,3}

¹*Institute for Theoretical Physics, University of California, Santa Barbara, CA, 93106*

²*Department of Physics, University of Houston, TX, 77204*

³*Department of Physics, Rutgers university, Piscataway, New Jersey, 07974*

(February 1, 2008)

By using mutual flux-attaching singular gauge transformations, we derive an effective action describing the zero temperature quantum phase transition from d-wave superconductor to underdoped regime. In this effective action, quantum fluctuation generated vortices couple to quasi-particles by a mutual statistical interaction with statistical angle θ , the vortices are also interacting with each other by long range interactions due to charge fluctuation. Neglecting the charge fluctuation, we find a fixed line characterized by θ and calculate the universal spinon, vortex and mutual Hall drag conductivities which continuously depend on θ . Implications for double layer quantum Hall systems are given. When incorporating the charge fluctuation, we emphasize the importance of keeping the periodicity of the mutual CS term at $\theta = \pm 1/2$ to get the correct critical behaviors and point out possible future directions. The connection with Z_2 gauge theory is discussed.

I. INTRODUCTION

In this paper, we are trying to study the nature of zero temperature quantum phase transition from d-wave superconductor at $x > x_c$ to the underdoped regime at $x < x_c$ of the high temperature superconductors (Fig.1).




Fig 1: The temperature (T) versus doping (x) diagram of high T_c cuprates.

It is well known that inside the superconductor phase $x > x_c$, there are low energy quasi-particle excitations near the four nodes of the Fermi surface, the quantum phase fluctuation of the Cooper pair condensates are suppressed, the positive and negative vortices are bound together. As the doping decreases from the right to x_c , the phase fluctuation increases. At the underdoped regime $x < x_c$, the quantum phase fluctuations are so strong that they generate free $hc/2e$ vortices, destroy the long range phase coherence of the d-wave superconductor, therefore the superconducting ground state. However, the local short-range pairing still exists, the low energy quasi-particles at the four nodes remain. This quantum phase transition is driven by the condensation of $hc/2e$ vortices. Near the transition, there are strong coupling between the quasi-particles and the phase windings of

$hc/2e$ vortices.

A lot of work has been done on a closely related problem where quasi-particles are coupled to the vortices generated by external magnetic field inside the superconducting state [1–6]. In the following, we will first review these work, then will extend the methods developed in these work to study this quantum phase transition where the quasi-particles are coupled to the quantum fluctuation generated vortices.

Employing semi-classical approximation, Volovik pointed out that the circulating supercurrents around vortices induce Doppler energy shift to the quasi-particle spectrum, which leads to a finite density of states at the nodes [1]. This effect (Volovik effect) has been employed to attempt to explain the experimental observations [7] of longitudinal thermal conductivity κ_{xx} in [2].

Starting from BCS Hamiltonian, Anderson [3] employed the first single-valued singular gauge transformation to study quasi-particle dynamics in the mixed state. Unfortunately, Anderson made an incorrect mean field approximation which violates the "Time-Reversal" symmetry [6], therefore leads to the incorrect conclusion that there is Landau level quantization of energy levels of the quasi-particle. Franz and Tesanovic (FT) employed a different single-valued singular gauge transformation to map the quasi-particle in a square vortex lattice state to Dirac fermion moving in an effective periodic scalar and vector potential with *zero* average and studied the quasi-particle spectrum numerically [4,5]. They did not see the signature of Landau level quantization in their numerical calculations

Unfortunately, in the vortex lattice state, the vector potential only provides a periodic potential instead of scattering quasi-particles, its effect on the energy spectrum and other physical quantities is not evident. This motivated Ye to study the quasi-particle transport in the disordered vortex state where the random AB phase scattering may show its important effects. In Ref. [6], Ye

observed that because infinite thin $hc/2e$ vortices do *not* break the T reversal symmetry, any correct mean field theory should respect this T symmetry. He showed exactly there is no Landau level quantization due to this T symmetry. He applied the singular transformation in the FT gauge to *disordered* vortex state and found that the long-range logarithmic interaction between vortices suppresses the fluctuation of superfluid velocity (scalar potential), but does *not* affect the fluctuation of the internal gauge field. Therefore the scalar field acquires a "mass" determined by the vortex density, but the gauge field remains "massless". The quasi-particle scattering from the "massless" internal gauge field dominates over those from the well-known "massive" Volovik effect and the non-magnetic scattering at sufficient high magnetic field. This dominant scattering mechanism is a purely quantum mechanical effects which was completely missed by all the previous semi-classical treatments [2]. In fact, it is responsible for the behaviors of both κ_{xx} and κ_{xy} in high magnetic field observed in the experiments [7].

When the vortices are generated by quantum fluctuations or thermal fluctuations, themselves are moving around, new physics may arise. From Newton's third law, intuitively, the vortices must also feel the counter-acting AB phase coming from the quasi-particles. Therefore there is a mutual AB phase scattering between vortices and quasi-particles. In this paper, we enlarge the *one* singular gauge transformation developed in [3,4,6] to *two* mutual flux-attaching singular gauge transformations. We then apply them to study quasi-particles coupled to moving vortices near the zero temperature quantum critical point. In canonical quantization, we perform two singular transformations which are dual to each other to quasi-particles and moving vortices respectively. Just like conventional singular gauge transformation leads to conventional Chern-Simon (CS) term, the two mutual singular gauge transformations lead to mutual CS term. This elegant mutual CS term precisely describe the mutual AB phase scattering between vortices and quasi-particles. Alternatively, in path-integral presentation, the effective action describes the quasi-particle moving in both vector and scalar potentials due to the phase fluctuations of quantum generated vortices. By a duality transformation presented in Refs. [8,9] to a vortex representation, the quantum fluctuation generated vortices couple to quasi-particles by a mutual CS term, the vortices are also interacting with each other by long range logarithmic interactions due to charge fluctuation described by a Maxwell term. By carefully considering the periodicity of this mutual CS term, it can be shown that the effective action is essentially equivalent to the dual vortex representation derived by Senthil and Fisher (SF) from the elegant Z_2 gauge theory [10] except the Volovik effect missing in the Z_2 gauge theory is also explicitly incorporated into the present approach. Neglecting the charge fluctuation first and replacing the Z_2 mu-

tual CS theory by the $U(1)$ mutual CS theory, we find a fixed line characterized by the mutual statistical angle θ and calculate the universal spinon, vortex and mutual Hall drag conductivities which continuously depend on θ . This transition can also be viewed as a simplest confinement and deconfinement transition. The properties of the confinement and deconfinement phases on the two sides of the critical point are also discussed. We stressed explicitly that the periodicity of Z_2 mutual CS gauge theory on the lattice is *not* preserved by the $U(1)$ mutual CS theory in the continuum. When the $U(1)$ charge fluctuation is taken into account, we tentatively still replace the Z_2 mutual CS theory by the $U(1)$ CS mutual theory and treat both $U(1)$ fluctuations on the same footing. We find the $U(1)$ charge fluctuation turns the fixed line into a fixed point where charge $\pm e$ and spin 1/2 Dirac-like quasi-particle and charge $2e$ Cooper pairs are asymptotically decoupled. Although the result is pathological, it does suggest the spinon and holon are confined into electrons and Cooper pairs when condensing $hc/2e$ vortices, in contrast to the condensing of double strength hc/e vortices discussed in [13,12] and reviewed in appendix C. Although we are unable to solve the critical behaviors of a combined Z_2 and $U(1)$ theory in this paper, we do stress that Z_2 periodicity should be treated correctly to understand the critical behaviors of a theory with both discrete Z_2 and continuous $U(1)$ gauge fields.

The paper is organized as the following. In the next section, we derive the effective action with both the mutual C-S interaction and the charge fluctuation, first in the path integral language, and then in canonical quantization representation. In Sec. III, we concentrate on the effect of the mutual C-S interaction and calculate the critical exponents and universal conductivities for general statistical angle θ . Implications for double layers Quantum Hall System are given. Some details are relegated to appendix A. In appendix C, we concentrate on the effect of charge fluctuation which is the only gauge fluctuation in double strength hc/e vortices. In Sec IV, we discuss the combined effect of mutual C-S interaction and charge fluctuation, we also stress the importance to keep the periodicity of Z_2 gauge field when considering a theory with both discrete Z_2 and continuous $U(1)$ gauge fields. In Sec. VI, we discuss the connection of our approach to the current fashion Z_2 gauge theory and point out the possible open problems. In appendix B, we apply the method developed in the main text to quenched random vortex array.

II. THE EFFECTIVE ACTION

Following the notation in Refs. [4,6], we define $d_{\uparrow} = c_{\uparrow}(x)$, $d_{\downarrow} = c_{\downarrow}^{\dagger}$ [18]. Because there are equal number of positive and negative vortices, we divide the positive vortices into two subsets ϕ_{p1} , ϕ_{p2} and negative vortices into

two subsets ϕ_{n1}, ϕ_{n2} . We define $\phi_A = \phi_{p1} + \phi_{n1}, \phi_B = \phi_{p2} + \phi_{n2}$ and introduce the spinon by performing a general singular unitary transformation $d = U d_s$:

$$H_s = U^{-1} H U, \quad U = \begin{pmatrix} e^{i\phi_A} & 0 \\ 0 & e^{-i\phi_B} \end{pmatrix} \quad (1)$$

where $\phi_A + \phi_B = \phi$ is the phase of the Cooper pair. The original transformation [3,4] is devised for static vortices. Here we extend it to moving vortices whose phase ϕ is also fluctuating, therefore, depends on both the space and time. The spinon is charge neutral. In Anderson's gauge, $\phi_A = 0, \phi_B = \phi$ or vice versa. In the former(latter), the spinon is electron-like (hole-like).

Expanding H_s around the node 1 where $\vec{p} = (p_F, 0)$, we obtain the linearized quasi-particle Lagrangian $\mathcal{L}_{qp}$ in the presence of the external gauge potential A_μ :

$$\mathcal{L}_{qp}^u = \psi_1^\dagger [(\partial_\tau - ia_\tau) + v_f(p_x - a_x)\tau^3 + v_2(p_y - a_y)\tau^1] \psi_1 + \psi_1^\dagger \psi_1 v_f v_x(\vec{r}) + i\psi_1^\dagger \tau^3 \psi_1 v_\tau(\vec{r}) + (1 \rightarrow 2, x \rightarrow y) \quad (2)$$

where $v_\mu = \frac{\hbar}{2} \partial_\mu \phi - \frac{e}{c} A_\mu$ is the gauge-invariant superfluid *momentum*, it acts as a scalar scattering potential, $a_\mu = \frac{1}{2} \partial_\mu (\phi_A - \phi_B)$ is the AB gauge field due to the phase winding of vortices [14]. Note that the external gauge potential only appear explicitly in the superfluid momentum v_μ . Because there are equal number of positive and negative vortices in both ϕ_A and ϕ_B , on the average, the vanishing of v_μ and a_μ is automatically ensured, in this case, the Anderson gauge turns out to be more convenient than FT gauge. In the following, we use the electron-like Anderson gauge where $a_\mu = \frac{1}{2} \partial_\mu \phi$ [17].

We get the corresponding expression at node $\bar{1}$ and $\bar{2}$ by changing $v_f \rightarrow -v_f, v_2 \rightarrow -v_2$ in the above Eq.

$$\mathcal{L}_{qp}^l = \psi_{\bar{1}}^\dagger [(\partial_\tau - ia_\tau) - v_f(p_x - a_x)\tau^3 - v_2(p_y - a_y)\tau^1] \psi_{\bar{1}} - \psi_{\bar{1}}^\dagger \psi_{\bar{1}} v_f v_x(\vec{r}) + i\psi_{\bar{1}}^\dagger \tau^3 \psi_{\bar{1}} v_\tau(\vec{r}) + (\bar{1} \rightarrow \bar{2}, x \rightarrow y) \quad (3)$$

Performing a P-H transformation $\tilde{\psi}_{1\alpha} = \epsilon_{\alpha\beta} \psi_{\bar{1}\beta}^\dagger$ and the corresponding expression at node $\bar{2}$, the above Eq. becomes:

$$\mathcal{L}_{qp}^l = \tilde{\psi}_1^\dagger [(\partial_\tau + ia_\tau) + v_f(p_x + a_x)\tau^3 + v_2(p_y + a_y)\tau^1] \tilde{\psi}_1 + \tilde{\psi}_1^\dagger \tilde{\psi}_1 v_f v_x(\vec{r}) + i\tilde{\psi}_1^\dagger \tau^3 \tilde{\psi}_1 v_\tau(\vec{r}) + (1 \rightarrow 2, x \rightarrow y) \quad (4)$$

In order to make the final expressions explicitly $SU(2)$ invariant, we perform the singular gauge transformation $\psi_{12\alpha} = e^{-i(\phi_A - \phi_B)} \tilde{\psi}_{1\alpha}$ and the corresponding transformation at node $\bar{2}$, then $a_\mu \rightarrow -a_\mu$, Eq.4 takes the same form as Eq.2 [15]. Adding the two equations leads to:

$$\mathcal{L}_{qp} = \psi_{1a}^\dagger [(\partial_\tau - ia_\tau) + v_f(p_x - a_x)\tau^3 + v_2(p_y - a_y)\tau^1] \psi_{1a} + \psi_{1a}^\dagger \psi_{1a} v_f v_x(\vec{r}) + i\psi_{1a}^\dagger \tau^3 \psi_{1a} v_\tau(\vec{r}) + (1 \rightarrow 2, x \rightarrow y) \quad (5)$$

where $a = 1, 2$ is the spin indices. τ^s matrices are acting on *particle-hole* space. As intended, the above Eq.

is explicitly $SU(2)$ spin invariant. Equivalently, we can start with the explicitly spin $SU(2)$ invariant approach advocated in Ref. [13] and perform the singular gauge transformation in the *p-h* space.

$\mathcal{L}_{qp}$ enjoys gauge symmetry $U(1)_u \times U_s(1)$ (in fact, it is $U(1) \times Z_2$), the first being uniform and second being staggered gauge symmetry: Uniform (or external) $U_u(1)$ gauge symmetry

$$c_\alpha \rightarrow c_\alpha e^{i\chi}, \quad d_\alpha \rightarrow d_\alpha \\ \phi_A \rightarrow \phi_A + \chi, \quad \phi_B \rightarrow \phi_B + \chi \quad (6)$$

Under this uniform $U(1)$ transformation, the corresponding fields transform as:

$$\phi \rightarrow \phi + 2\chi, \quad A_\alpha \rightarrow A_\alpha + \partial_\alpha \chi \\ v_\alpha \rightarrow v_\alpha, \quad a_\alpha \rightarrow a_\alpha \quad (7)$$

d, v_α, a_α all are invariant under this external $U(1)$ transformation. Therefore the spinon d_α is charge *neutral* to the external magnetic field.

Staggered (or internal) $U_s(1)$ gauge symmetry

$$c_\alpha \rightarrow c_\alpha, \quad d_\alpha \rightarrow d_\alpha e^{-i\chi} \\ \phi_A \rightarrow \phi_A + \chi, \quad \phi_B \rightarrow \phi_B - \chi \quad (8)$$

Under this internal $U(1)$ transformation, the corresponding fields transform as:

$$\phi \rightarrow \phi, \quad A_\alpha \rightarrow A_\alpha \\ v_\alpha \rightarrow v_\alpha, \quad a_\alpha \rightarrow a_\alpha + \partial_\alpha \chi \quad (9)$$

Although the spinon d_α is charge neutral to the external magnetic field, it carries charge 1 to the internal gauge field a_α .

It is easy to realize that $U_u(1)$ acts only on the boson sector, since the spinon is charge neutral, $U_s(1)$ acts only on the fermion sector. In fact, $U_s(1)$ should be a discrete local Z_2 symmetry [10], because up and down $\hbar c/2e$ vortices are equivalent and do not break T symmetry [6].

The phase fluctuation is simply 2 + 1 dimensional X-Y model:

$$\mathcal{L}_{ph} = \frac{K}{2} v_\mu^2 = \frac{K}{2} (\partial_\mu \phi - 2A_\mu)^2 \quad (10)$$

After absorbing the scalar potential scattering part (Volovik term) into $\mathcal{L}_{ph}$, we can write the total Lagrangian $\mathcal{L} = \mathcal{L}_{qp} + \mathcal{L}_{ph}$ as:

$$\mathcal{L} = \psi_{1a}^\dagger [(\partial_\tau - ia_\tau) + v_f(p_x - a_x)\tau^3 + v_2(p_y - a_y)\tau^1] \psi_{1a} + (1 \rightarrow 2, x \rightarrow y) + \frac{K}{2} (\partial_\mu \phi - A_\mu^{eff})^2 \quad (11)$$

where $A_\mu^{eff} = 2A_\mu - K^{-1} J_\mu$ and the quasi-particle electric current is : $J_0 = \psi_j^\dagger \tau^3 \psi_j, J_x = v_f \psi_1^\dagger \psi_1, J_y = v_f \psi_2^\dagger \psi_2$.

From Eq.11, it is easy to identify the two conserved Noether currents: spinon current and electric current.

The spinon current is given by:

$$\begin{aligned} j_0^s &= \psi_1^\dagger(x)\psi_1(x) + \psi_2^\dagger(x)\psi_2(x) \\ j_x^s &= \psi_1^\dagger(x)v_F\tau^3\psi_1(x) + \psi_2^\dagger(x)v_2\tau^1\psi_2(x) \\ j_y^s &= \psi_1^\dagger(x)v_2\tau^1\psi_1(x) + \psi_2^\dagger(x)v_F\tau^3\psi_2(x) \end{aligned} \quad (12)$$

Obviously the spinon current only comes from quasi-particle. In principle, the spinon current is not conserved due to scatterings between different nodes which lead to anomalous terms not included in Eq.11. However, the inter-node scatterings involve large momenta transfer $\vec{K}_i - \vec{K}_j$ for $i \neq j$, so we neglect them due to momentum conservation in the long wave-length limit of the phase fluctuation [16]. The exact conserved *spin current* $\vec{j}_\mu^S$ is with $\vec{\sigma}/2$ inserted in the above spinon currents.

The electric current is given by:

$$j_\mu^e = -\frac{\partial \mathcal{L}}{\partial A_\mu^{eff}} = K(\partial_\mu \phi - A_\mu^{eff}) = K(\partial_\mu \phi - 2A_\mu) + J_\mu \quad (13)$$

Where the first part coming from Cooper pair and the second from the quasi-particle. Although they are not separately conserved, their sum is.

The Noether current due to the the symmetry under $\phi \rightarrow \phi + \chi$ can be written as:

$$j_\mu^t = \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi)} = K(\partial_\mu \phi - A_\mu^{eff}) - \frac{1}{2}j_\mu^s = j_\mu^e - \frac{1}{2}j_\mu^s \quad (14)$$

It is a combination of electric and spinon currents, therefore also conserved.

Following Refs. [8,9], we perform a duality transformation to Eq.11

$$\begin{aligned} &\frac{K}{2}(\partial_\mu \phi - A_\mu^{eff})^2 - i\frac{1}{2}\partial_\mu \phi j_\mu^s \\ &= ij_{e\mu}(\partial_\mu \phi - A_\mu^{eff}) + \frac{1}{2K}j_{e\mu}^2 - i\frac{1}{2}\partial_\mu \phi j_\mu^s \\ &= ij_\mu^t \partial_\mu \theta + ij_\mu^t \partial_\mu \phi - ij_\mu^e A_\mu^{eff} + \frac{1}{2K}j_{e\mu}^2 \end{aligned} \quad (15)$$

Where we have separated topological trivial spin-wave part and topological non-trivial vortex parts.

Integrating out the spin-wave part, we get the conservation equation for the total current $j_\mu^t = j_\mu^e - \frac{1}{2}j_\mu^s$. In fact, as shown in the previous paragraphs, j_μ^e and j_μ^s are separately conserved. Therefore we can introduce spin and electric gauge fields by:

$$\begin{aligned} j_\mu^s &= \epsilon_{\mu\nu\lambda} \partial_\nu a_\lambda^s \\ j_\mu^e &= \epsilon_{\mu\nu\lambda} \partial_\nu a_\lambda^e \end{aligned} \quad (16)$$

We can also define the vortex current:

$$\begin{aligned} j_\mu^v &= \frac{1}{2\pi} \epsilon_{\mu\nu\lambda} \partial_\nu \partial_\lambda \phi \\ &= \epsilon_{\mu\nu\lambda} \partial_\nu a_\lambda^v \end{aligned} \quad (17)$$

where $a_\mu^v = \partial_\mu \phi / 2\pi$ is the vortex gauge field.

Substituting the above expressions into Eq.15, we reach:

$$\begin{aligned} &\frac{1}{4K} f_{e\mu\nu}^2 + i\partial_\mu \phi j_\mu^t - iA_\mu^{eff} \epsilon_{\mu\nu\lambda} \partial_\nu a_\lambda^e \\ &= \frac{1}{4K} f_{e\mu\nu}^2 + ia_\mu^t j_\mu^v - iA_\mu^{eff} \epsilon_{\mu\nu\lambda} \partial_\nu a_\lambda^e \end{aligned} \quad (18)$$

where $a_\mu^t = a_\mu^e - \frac{1}{2}a_\mu^s$ is the total gauge field felt by the moving vortices.

Using Φ for the vortex operator and adding the quasi-particle part, we get the following effective action:

$$\begin{aligned} \mathcal{L} &= \psi_1^\dagger[(\partial_\tau - ia_\tau^\psi) + v_f(p_x - a_x^\psi)\tau^3 + v_2(p_y - a_y^\psi)\tau^1]\psi_1 \\ &+ (1 \rightarrow 2, x \rightarrow y) \\ &+ |(\partial_\mu - ia_\mu^\Phi - ia_\mu) \Phi|^2 + V(|\Phi|) + \frac{i}{2\pi\theta} a_\mu^\psi \epsilon_{\mu\nu\lambda} \partial_\nu a_\lambda^\Phi \\ &+ \frac{1}{4} f_{\mu\nu}^2 - iA_\mu^{eff} \epsilon_{\mu\nu\lambda} \partial_\nu a_\lambda - \mu \epsilon_{ij} \partial_i a_j \end{aligned} \quad (19)$$

where $V(|\Phi|) = m_\Phi^2 |\Phi|^2 + g_\Phi |\Phi|^4 + \dots$ stands for the short range interaction between the vortices. The last term is due to Berry phase in the boson representation [9] which can be absorbed into A_μ^{eff} by redefining $A_\mu^{eff} \rightarrow A_\mu^{eff} + i\mu\delta_{\mu 0}$, it acts like an external magnetic field in the $\hat{z}$ (namely $\mu = 0$) direction. There are one species of vortex and $N = 4$ species of Dirac fermion(2 spin components at 2 nodes at the upper plane), the mutual statistical angle $\theta = \pm 1/2$. We also changed the notation by setting $a^\psi = a^v, a^\Phi = a^s, a^e = a$.

Although we derived the above equation in the electron-like Anderson gauge, in fact, it holds in *any* gauge. Eq.19 again enjoys the gauge symmetry $U(1) \times U(1)$, the first only acting on the vortex sector is the electric $U(1)$ gauge field, the second on both the vortex and fermion sectors is the $U(1)$ mutual CS gauge field. The mutual Chern-Simon term enforces the constraints: $2\pi\theta j_\mu^s = \epsilon_{\mu\nu\lambda} \partial_\nu a_\lambda^\Phi, 2\pi\theta j_\mu^v = \epsilon_{\mu\nu\lambda} \partial_\nu a_\lambda^\psi$. Physically, it means that when a quasi-particle encircles a vortex, it picks up a phase $2\pi\theta$. Equivalently when a vortex moves around a quasi-particle, it also picks up a phase $2\pi\theta$. Although the conventional C-S term breaks T symmetry and has periodicity under $\theta \rightarrow \theta + 2$ [19–23], the mutual C-S term does *not* break T symmetry and has periodicity under $\theta \rightarrow \theta + 1$. For example, $\theta = -1/2$ is equivalent to $\theta = 1/2$. In fact, this mutual CS term at $\theta = 1/2$ should be the same as the Z_2 mutual CS term on the lattice discussed in Ref. [10]. Therefore strictly speaking a^ψ and a^Φ are Z_2 gauge fields, only a_μ is a $U(1)$ gauge field. The Doppler-shifted term (Volovik effect) is encoded in the last term in Eq.19. This effect was *not* taken into account in the Z_2 gauge theory [10].

In canonical quantization language, it is very instructive to compare the well-known singular gauge transformation leading to composite fermion in $\nu = 1/2$ system by Halperin, Lee and Read (HLR) [19] to the singular gauge transformations performed in the spinon-vortex system in this paper. The crucial difference is that in the former, we attach electron's own θ flux to itself ($\theta = 2$ in $\nu = 1/2$ system to keep fermion statistics intact), the constraint $\nabla \times \vec{a}(\vec{r}) = 2\pi\theta\rho(\vec{r})$ in Coulomb gauge $\partial_i a_i = 0$ leads to composite fermion ψ_c coupled to conventional CS term which breaks T symmetry, has periodicity under $\theta \rightarrow \theta + 2$; however, in the latter, we attach vortex's $\theta = 1/2$ flux to quasi-particle or *vice versa*, the two constraints $\nabla \times \vec{a}^\psi(\vec{r}) = 2\pi\theta j_0^v(\vec{r})$, $\nabla \times \vec{a}^\Phi(\vec{r}) = 2\pi\theta j_0^s(\vec{r})$ in Coulomb gauges $\partial_i a_i^\psi = \partial_i a_i^\Phi = 0$ naturally leads to quasi-particles and vortices are coupled by mutual CS term in addition to the electrical gauge field which mediates the logarithmic interaction between vortices. The mutual CS term does not break T symmetry, has periodicity under $\theta \rightarrow \theta + 1$.

It is important to point out that when a vortex is moving around a closed loop, it pick up two phases, one is Z_2 phase due to the spinon current, described by the mutual C-S term, another is $U(1)$ phase due to the total electric charge current described by the Maxwell term. In the following section, we will neglect the electric charge fluctuation and concentrate the mutual statistics term. In Sec. IV, we will study the effect of charge fluctuation described by the Maxwell term. For completeness, we relegate the review of the effect of charge fluctuation to appendix B.

For simplicity, we take the relativistic form for both fermion and boson, because the anisotropies in Eq.19 are expected to be irrelevant near the zero temperature QCP [11].

III. THE EFFECT OF MUTUAL STATISTICS

In this section, we neglect the charge fluctuation, namely setting $a_\mu = 0$ in Eq.19. The charge fluctuation can be suppressed by condensing hc/e vortices reviewed in appendix C. We could add the kinetic and potential terms for the hc/e vortex operator Φ_2 to Eq.19:

$$\mathcal{L}_{\Phi_2} = |(\partial_\mu - i2a_\mu)\Phi_2|^2 + V(|\Phi_2|) \quad (20)$$

As shown in appendix C, there is no mutual statistical interaction between spinon and hc/e vortices Φ_2 . The long range logarithmic interaction between $hc/2e$ vortices Φ and hc/e vortices Φ_2 is mediated by the electrical gauge field a_μ . Condensing $\langle \Phi_2 \rangle = \Phi_{20}$ will generate a mass term $\frac{\Phi_{20}^2}{2}(a_\mu)_i^2$ which dominates over the Maxwell term. Integrating out a_μ leads to

$$\frac{1}{\Phi_{20}^2} \left[\frac{1}{4} (f_{\mu\nu}^{Aeff})^2 - iA_\mu^{eff} \epsilon_{\mu\nu\lambda} \partial_\nu j_\lambda^v + (j_\mu^v)_i^2 \right] \quad (21)$$

All the generated terms only renormalize the short range interactions already included in $V(|\Phi|)$. The model is also closely related to the Hall drag problem in double layer Quantum Hall system [26], therefore the model itself is interesting on its own right and deserves detailed investigation.

A. Quantum Critical point

In order to calculate conductivities, we add two source fields A_μ^ψ and A_μ^Φ for the quasi-particles and vortex respectively:

$$\mathcal{L} = \psi_a^\dagger \gamma_\mu (\partial_\mu - ia_\mu^\psi - iA_\mu^\psi) \psi_a + |(\partial_\mu - ia_\mu^\Phi - iA_\mu^\Phi)\Phi|^2 + V(|\Phi|) + \frac{i}{2\pi\theta} a_\mu^\psi \epsilon_{\mu\nu\lambda} \partial_\nu a_\lambda^\Phi \quad (22)$$

where $a = 1, 2, 3, 4$ stands for $N = 4$ species of Dirac fermion. In fact, two Maxwell terms for a^ψ and a^Φ can be added to the above equation, but they are subleading to the mutual CS term in the low energy limit. It is expected that there is no periodicity under $\theta \rightarrow \theta + 1$ in the continuum limit.

If the theory is properly regularized on the lattice, it should have the periodicity under $\theta \rightarrow \theta + 1$, for example, $\theta = -1/2$ should be equivalent to $\theta = 1/2$. In fact, $\theta = 1/2$ mutual $U(1)$ CS theory could be the same as the Z_2 mutual CS theory developed in [10].

The RG calculation in Refs. [20–22] can be used to show that θ is exactly marginal, therefore there is a line of fixed points determined by the mutual statistical angle θ . In order to calculate the *spin conductivity* along this fixed line, a source field could be introduced to couple to the spin current $\vec{j}_\mu^S = \psi_\alpha^\dagger \gamma_\mu (\vec{\sigma})_{\alpha\beta} \psi_\beta$. Similar calculations follow.

Integrating out both fermion and boson leads to:

$$\begin{aligned} \mathcal{L} = & -\frac{1}{2} a_\mu^\psi(-k) \Pi_{\mu\nu}^\psi(k) a_\nu^\psi(k) - \frac{1}{2} a_\mu^\psi(-k) \Pi_{\mu\nu}^{\psi\Phi}(k) a_\nu^\Phi(k) \\ & - \frac{1}{2} a_\mu^\Phi(-k) \Pi_{\mu\nu}^\Phi(k) a_\nu^\Phi(k) - \frac{1}{2} a_\mu^\Phi(-k) \Pi_{\mu\nu}^{\Phi\psi}(k) a_\nu^\psi(k) \\ & - \frac{1}{2\theta} (a_\mu^\psi(-k) - A_\mu^\psi(-k)) \epsilon_{\mu\nu\lambda} k_\lambda (a_\nu^\Phi(k) - A_\nu^\Phi(k)) \\ & - \frac{1}{2\theta} (a_\mu^\Phi(-k) - A_\mu^\Phi(-k)) \epsilon_{\mu\nu\lambda} k_\lambda (a_\nu^\psi(k) - A_\nu^\psi(k)) \end{aligned} \quad (23)$$

where the exact forms of Π^s are dictated by gauge invariance and Furry's theorem :

$$\begin{aligned} \Pi_{\mu\nu}^\psi(k) &= \Pi_1(k) k (\delta_{\mu\nu} - \frac{k_\mu k_\nu}{k^2}) \\ \Pi_{\mu\nu}^{\psi\Phi}(k) &= \Pi_{\mu\nu}^{\Phi\psi}(k) = \Pi_2(k) \epsilon_{\mu\nu\lambda} k_\lambda \\ \Pi_{\mu\nu}^\Phi(k) &= \Pi_3(k) k (\delta_{\mu\nu} - \frac{k_\mu k_\nu}{k^2}) \end{aligned} \quad (24)$$

Where Π_1, Π_2, Π_3 are the polarizations for $a^\psi a^\psi, a^\psi a^\Phi, a^\Phi a^\Phi$.

If we are only interested in DC conductivities, for simplicity, we can put $\vec{k} = 0$, Eq.23 becomes (for the most general form, see appendix A):

$$\begin{aligned} \mathcal{L} = & -\frac{1}{2}a_i^\psi(-\omega_n)\Pi_1|\omega_n|a_i^\psi(\omega_n) - \frac{1}{2}a_i^\psi(-\omega_n)\Pi_2\epsilon_{ij}\omega_n a_j^\Phi(\omega_n) \\ & - \frac{1}{2}a_i^\Phi(-\omega_n)\Pi_3|\omega_n|a_i^\Phi(\omega_n) - \frac{1}{2}a_i^\Phi(-\omega_n)\Pi_2\epsilon_{ij}\omega_n a_j^\psi(\omega_n) \\ & - \frac{1}{2\theta}(a_i^\psi(-\omega_n) - A_i^\psi(-\omega_n))\epsilon_{ij}\omega_n(a_j^\Phi(\omega_n) - A_j^\Phi(\omega_n)) \\ & - \frac{1}{2\theta}(a_i^\Phi(-\omega_n) - A_i^\Phi(-\omega_n))\epsilon_{ij}\omega_n(a_j^\psi(\omega_n) - A_j^\psi(\omega_n)) \end{aligned} \quad (25)$$

If we define $\tilde{a}_i^\Phi(\omega_n) = \epsilon_{ij}a_j^\Phi(\omega_n)$, $\tilde{A}_i^\Phi(\omega_n) = \epsilon_{ij}A_j^\Phi(\omega_n)$, the above equation becomes diagonal in the spatial indices $i = 1, 2$. Finally, integrating out $a_i^\psi, \tilde{a}_i^\Phi$ leads to

$$\mathcal{L} = -\frac{1}{2}(A_i^\psi, \tilde{A}_i^\Phi) \begin{pmatrix} \sigma^\psi & \sigma^H \\ \sigma^H & \sigma^\Phi \end{pmatrix} \begin{pmatrix} A_i^\psi \\ \tilde{A}_i^\Phi \end{pmatrix} \quad (26)$$

Where spinon, vortex and mutual Hall drag conductivities are:

$$\begin{aligned} \sigma^\psi &= \left(\frac{1}{\theta}\right)^2 \frac{\Pi_1}{\Pi_1\Pi_3 + (1/\theta - \Pi_2)^2} \\ \sigma^\Phi &= \left(\frac{1}{\theta}\right)^2 \frac{\Pi_3}{\Pi_1\Pi_3 + (1/\theta - \Pi_2)^2} \\ \sigma^H &= \left(\frac{1}{\theta}\right)^2 \frac{\Pi_2 - \theta(\Pi_1\Pi_3 + \Pi_2^2)}{\Pi_1\Pi_3 + (1/\theta - \Pi_2)^2} \end{aligned} \quad (27)$$

In fact, all the three conductivities can be written in the elegant connection formula:

$$\rho_{ij} = (\rho_{FB})_{ij} - \theta\epsilon_{ij} \quad (28)$$

with the conductivity tensor of fermion and boson given by

$$\sigma_{FB} = \begin{pmatrix} \Pi_1 & -\Pi_2 \\ \Pi_2 & \Pi_3 \end{pmatrix} \quad (29)$$

Although its form is similar to the conventional connection formulas discussed in [19–21], the physical interpretations of the conductivities are quite different (see the following).

When the vortices are generated by external magnetic field and pinned by impurities as discussed in [6], the total conductivity is the same as the fermion conductivity because the static vortex do not contribute. As explained in appendix B, the static vortices only feel the electric gauge field a_μ , but *not* the statistical gauge field a_μ^Φ .

The above expressions are exact, but Π_1, Π_2, Π_3 can only be calculated perturbatively in the coupling constant $g^2 = 2\pi\theta$. The renormalized propagators for a^ψ and a^Φ can be found from the following Feymann diagrams:




Fig 2: The renormalized propagators for a^ψ (thick wiggly line) and a^Φ (thick dashed line). The thin wiggly line stands for a^ψ , the dashed thin line for a^Φ , the thin solid line for fermion propagators, the thick solid line for the boson propagators.

By using the bare propagators $\langle a_\mu^\psi a_\nu^\Phi \rangle = -\epsilon_{\mu\nu\lambda}k_\lambda/k^2$ and the bare fermion and boson loop results $\Pi_{\mu\nu}^{f0} = \Pi_{\mu\nu}^{b0} = -\frac{g^2}{16}k(\delta_{\mu\nu} - k_\mu k_\nu/k^2)$, we can find easily the renormalized a^ψ and a^Φ propagators $G_{\mu\nu}^\psi = G_{\mu\nu}^\Phi = -\frac{g^2}{16} \frac{1}{k}(\delta_{\mu\nu} - k_\mu k_\nu/k^2)$. In contrast to the conventional CS theory studied in [20,21], the propagators are *even* in k , this is because the theory respects T symmetry. On the other hand, in contrast to the Maxwell propagators, they behave as $1/k$ instead of $1/k^2$.

The three loop diagrams for Π_1 are given by:




Fig 3: The three loop diagrams of Π_1 . The thick wiggly line stands for the renormalized propagators of a^ψ , the thin solid line stands for the fermion propagator. The one loop diagram is not shown.

By using $G_{\mu\nu}^\psi$ and extracting the symmetric part of the gauge propagator in the large N results in Refs. [20,21], we are able to calculate the above three loop diagrams. Furry's theorem can be used to eliminate large number of null diagrams. We get the following series:

$$\Pi_1 = N \frac{\pi}{8} \left(1 + \frac{3}{16} \frac{g^4}{(2\pi)^2} + g^8 + \dots \right) \quad (30)$$

where $N = 4$ is due to the sum over 4 species of Dirac fermions.

The three loop diagrams for Π_3 are given by:




Fig 4: The three loop diagrams of Π_3 . The thick dashed line stands for the renormalized propagator of a^Φ , the thick solid line stands for the boson propagator. The one loop diagram is not shown.

For bosons, only one loop result is known:

$$\Pi_3 = N \frac{\pi}{8} (1 + g^4 + g^8 + \dots) \quad (31)$$

Note that both Π_1 and Π_3 are even functions of θ .

From Furry's theorem, one of the first non-vanishing diagram for Π_2 is:




Fig 5: One of the four loop diagrams of Π_2 . the thin solid line stands for the fermion propagator, the thick solid line for the boson propagator. All the other four loop diagrams can be obtained by shuffling the positions of the three bare propagator lines.

The series is $\Pi_2 = N(g^6 + g^{10} + \dots)$ which is an *odd* function of θ .

From Eq.27, σ^ψ, σ^Φ are even functions of θ , but the mutual Hall drag conductivity is an *odd* function of θ . These are expected from P-H transformation. Under the P-H transformation of the vortex operator $\Phi \rightarrow \Phi^\dagger$ in Eq.22, it can be shown that A_μ^Φ, θ is equivalent to $-A_\mu^\Phi, -\theta$. From Eq.26, we reach the same conclusions. Specifically, σ^H takes opposite values for $\theta = \pm 1/2$, the periodicity under $\theta \rightarrow \theta + 1$ is not preserved. Experimentally, the Hall drag conductivity can be detected by measuring the transverse voltage drop (or transverse temperature drop for thermal conductivity) of spinons due to the longitudinal driving of vortices. The Hall drag conductivity in double layer Quantum Hall systems has been investigated by several authors [26]. In double layer systems, the electrons in different layers are treated as two different species. There is a mutual CS interaction between the two species (both are fermions) which is directly responsible for this Hall drag conductivity, although the Coulomb interaction between the two species is responsible for the Coulomb (longitudinal) drag [27]. This example shows that *no* external magnetic field is needed to produce a Hall effect ! However, $\theta = \pm 1/2$ are very special points, $U(1)$ mutual CS term reduces to Z_2 field, there should be *no* Hall drag conductivity just like $\alpha = \pm 1/2$ vortex leads to *no* Hall conductivity.

Again, by using the large N result of [20,21], we find the anomalous dimensions of the fermion and vortex to two loops:




Fig 6: The two loop diagrams for the self energies of spinon (a) and vortex (b). The renormalized operators are used.

The results are:

$$\eta_\psi = -\frac{g^4}{48\pi^2}$$

$$\eta_\Phi = \eta_{XY} - \frac{g^4}{12\pi^2}N \quad (32)$$

where $\eta_{XY} \sim 0.038$ is the anomalous dimension for the 3d XY model [24].

The correlation length exponent can also be calculated to two loops by the insertion of the operator $\Phi^\dagger\Phi$:




Fig 7: The cross stands for the operator insertion of $\Phi^\dagger\Phi$.

The result is:

$$\nu = \nu_{XY} - \frac{g^4}{12\pi^2}N \quad (33)$$

where $\nu_{XY} \sim 0.672$ is correlation length exponent for the 3d XY model [24].

It is instructive to go to dual representation of Eq.22, namely go to the boson representation:

$$\mathcal{L} = \psi_a^\dagger \gamma_\mu (\partial_\mu - ia_\mu^\psi - iA_\mu^\psi) \psi_a + |(\partial_\mu - ia_\mu^\phi)\phi|^2 + V(|\phi|) + \frac{1}{4}(f_{\mu\nu}^\phi)^2 + ia_\mu^\Phi \epsilon_{\mu\nu\lambda} \partial_\nu (a_\lambda^\phi - a_\lambda^\psi/\theta) - iA_\mu^\Phi \epsilon_{\mu\nu\lambda} \partial_\nu a_\lambda^\phi \quad (34)$$

where $V(|\phi|) = m_\phi^2 |\phi|^2 + g_\phi |\phi|^4 + \dots$,

Integrating out a_μ^Φ leads to the constraint up to a pure gauge:

$$a_\mu^\psi = \theta a_\mu^\phi \quad (35)$$

Substituting the above constraint to Eq.34 and setting $a_\mu^\phi = a_\mu$, we find:

$$\mathcal{L} = \psi_a^\dagger \gamma_\mu (\partial_\mu - i\theta a_\mu - iA_\mu^\psi) \psi_a + \frac{1}{4}f_{\mu\nu}^2 + |(\partial_\mu - ia_\mu)\phi|^2 + V(|\phi|) - iA_\mu^\Phi \epsilon_{\mu\nu\lambda} \partial_\nu a_\lambda \quad (36)$$

The above Eq. indicates that fermions and bosons are coupled to the same gauge field whose dynamics is described by Maxwell term instead of C-S term. This Eq. is simply 2 + 1 dimensional combination of spinor QED and scalar QED. The RG analysis at $4 - \epsilon$ by dimensional regularization is possible, because the marginal dimension of all the relevant couplings are 4. However a RG analysis directly at 2 + 1 dimension is formidable, the physical meaning and the exact marginality of θ is obscure in the boson representation. However they are evident in the dual vortex representation Eq.22. It is evident that there is no periodicity under $\theta \rightarrow \theta + 1$ in Eq.36.

If we perform duality transformation again on Eq.36 to go to the vortex representation, then we recover Eq.19 upon neglecting the two Maxwell terms relative to the mutual C-S term.

In the next subsection, setting the two source terms vanishing, we look at the properties of the different phases on the two sides of this quantum critical point.

B. Disorder and superconducting phases

In the disordered phase, the vortex condense $\langle \Phi \rangle = \Phi_0$ which generates a mass term for a_μ^Φ in Eq.22

$$\frac{\Phi_0^2}{2}(a_\mu^\Phi)_t^2 \quad (37)$$

where the subscript t means transverse projection.

Integrating out the massive a_μ^Φ leads to a Maxwell term for a_μ^ψ :

$$\mathcal{L} = \psi_a^\dagger \gamma_\mu (\partial_\mu - i a_\mu^\psi) \psi_a + \frac{1}{4\Phi_0^2} (f_{\mu\nu}^\psi)^2 \quad (38)$$

This is simply 2 + 1 dimensional spinor QED which was studied by large N expansion in [25]. In fact, we reach the same description from the boson representation Eq.36. Because in the disordered phase, the boson ϕ is massive, therefore can be integrated out, it generates the Maxwell term $\frac{1}{4m_\phi} f_{\mu\nu}^2$ which dominates over the existing *non-critical* Maxwell term. We reach Eq.38 after identifying $m_\phi \sim \Phi_0^2$.

In the superconductor phase, the vortex Φ is massive, therefore can be integrated out, it leads to the old Maxwell term $\frac{1}{4m_\Phi} (f_{\mu\nu}^\Phi)^2$. Integrating out a^Φ generates a mass term:

$$\mathcal{L} = \psi_a^\dagger \gamma_\mu (\partial_\mu - i a_\mu^\psi) \psi_a + \frac{m_\Phi}{2} (a_\mu^\psi)_t^2 \quad (39)$$

where m_Φ is the mass of the vortex.

The Dirac fermions become free. In fact, we reach the same description from the boson representation Eq.36. Because in superconductor phase, the boson ϕ condense, therefore generates a mass term $\frac{\phi_0^2}{2} (a_\mu)^2$ which renders the Maxwell term ineffective. We reach the same conclusion from both sides by identifying $m_\Phi \sim \phi_0^2$.

In short, in the disordered phase, the system is described by spinor QED Eq.38; in the superconductor side, by free Dirac fermion. We can view the transition as the simplest confinement-deconfinement transition. In the confined (disordered) phase, the boson and fermion are confined together by the fluctuating gauge field. In the deconfined (superconductor) phase, the boson condensed, the fermion becomes free. There is a line of fixed point governed by the mutual statistical angle θ separating the two phases.

IV. THE EFFECT OF CHARGE FLUCTUATION

In this section, we try to investigate the effect of charge fluctuation on the fixed line characterized by the statistical angle θ discussed in the last section. In the last section, we pointed out the periodicity of Z_2 gauge CS term on the *lattice* is not preserved by $U(1)$ gauge CS

term in the *continuum*. In this section, we try to consider the combined effects of Z_2 mutual statistical gauge fluctuation and $U(1)$ electrical gauge fluctuation. After replacing the Z_2 discrete gauge theory by a $U(1)$ mutual CS theory, we treat $U(1)$ mutual CS theory and $U(1)$ electrical gauge theory on the equal footing. Although the replacement leads to somewhat misleading conclusions, the heuristics derivation suggests that the condensation of $hc/2e$ vortex condensation indeed leads to the confinement of spinon and chargon into Cooper pair and electron, in contrast to the condensation of hc/e vortex. We do not intend to understand the critical behaviors of a combined Z_2 and $U(1)$ theory in this paper. We simply stress that Z_2 periodicity should be treated correctly to understand the critical behaviors of a theory with both discrete Z_2 and continuous $U(1)$ gauge fields.

A. Quantum critical point

Putting $a_\mu^\Phi \rightarrow a_\mu^\Phi - a_\mu$ in Eq.19, we get

$$\mathcal{L} = \psi_a^\dagger \gamma_\mu (\partial_\mu - i a_\mu^\psi) \psi_a + |(\partial_\mu - i a_\mu^\Phi) \Phi|^2 + V(|\Phi|) + \frac{i}{2\pi\theta} a_\mu^\psi \epsilon_{\mu\nu\lambda} \partial_\nu a_\lambda^\Phi - i a_\mu^\psi \epsilon_{\mu\nu\lambda} \partial_\nu (a_\lambda^\psi / \theta + A_\lambda^{eff}) + \frac{1}{4} f_{\mu\nu}^2 \quad (40)$$

Integrating out the electric gauge field a_μ leads to:

$$\mathcal{L} = \psi_a^\dagger \gamma_\mu (\partial_\mu - i a_\mu^\psi) \psi_a + |(\partial_\mu - i a_\mu^\Phi) \Phi|^2 + V(|\Phi|) + \frac{i}{2\pi\theta} a_\mu^\psi \epsilon_{\mu\nu\lambda} \partial_\nu a_\lambda^\Phi + \frac{1}{2} (a_\mu^\psi / \theta + A_\mu^{eff})_t^2 \quad (41)$$

Comparing to Eq.22, it is easy to see that the charge fluctuation leads to a mass term for the gauge field a_μ^ψ . Shifting $a_\mu^\psi / \theta + A_\mu^{eff} \rightarrow a_\mu$ and adding the gauge fixing term $\frac{1}{2\alpha} (\partial_\mu a_\mu)^2$, we can integrate out the *massive* gauge field a_μ in Lorenz gauge $\alpha = 0$ and find:

$$\mathcal{L} = \psi_a^\dagger \gamma_\mu (\partial_\mu - i \theta A_\mu^{eff}) \psi_a + |(\partial_\mu - i a_\mu^\Phi) \Phi|^2 + V(|\Phi|) + \frac{1}{4} (f_{\mu\nu}^\Phi)^2 - i (A_\mu^{eff} + \theta j_\mu^s) \epsilon_{\mu\nu\lambda} \partial_\nu a_\lambda^\Phi + (j_\mu^s)_t^2 \quad (42)$$

Note the Maxwell term for a^Φ is generated by the integration over the massive a_μ .

Setting $A_\mu = 0$, integrating out the fermions only leads to higher derivative terms than the Maxwell term:

$$\mathcal{L} = |(\partial_\mu - i a_\mu^\Phi) \Phi|^2 + V(|\Phi|) + \frac{1}{4} (f_{\mu\nu}^\Phi)^2 + \dots \quad (43)$$

where $\dots$ means higher than second order derivatives. Therefore the vortex and fermion are asymptotically decoupled. It indicates that the charge fluctuation neglected in the last section destroy the fixed line characterized by θ . However, very different conclusions are reached in Ref. [11]. We believe that the authors failed to treat the charge gauge field fluctuation correctly.

Just like the last section, it is instructive to go to dual representation of Eq.41, namely go to the boson representation:

$$\begin{aligned}\mathcal{L} = & \psi_a^\dagger \gamma_\mu (\partial_\mu - i a_\mu^\psi) \psi_a + |(\partial_\mu - i a_\mu^\phi) \phi|^2 + V(|\phi|) \\ & + \frac{1}{4} (f_{\mu\nu}^\phi)^2 + i a_\mu^\Phi \epsilon_{\mu\nu\lambda} \partial_\nu (a_\lambda^\phi - a_\lambda^\psi / \theta) \\ & + \frac{1}{2} (a_\mu^\psi / \theta + A_\mu^{eff})^2\end{aligned}\quad (44)$$

Integrating out a_μ^Φ leads to the *same* constraint as Eq.35 up to a pure gauge:

$$a_\mu^\psi = \theta a_\mu^\phi \quad (45)$$

Substituting the above constraint to Eq.44 and setting $a_\mu^\phi = a_\mu$, we find:

$$\begin{aligned}\mathcal{L} = & \psi_a^\dagger \gamma_\mu (\partial_\mu - i \theta a_\mu) \psi_a + |(\partial_\mu - i a_\mu) \phi|^2 + V(|\phi|) \\ & + \frac{1}{4} f_{\mu\nu}^2 + \frac{1}{2} (a_\mu + A_\mu^{eff})^2\end{aligned}\quad (46)$$

Comparing to Eq.36, the only difference is that the gauge field acquires a mass due to the charge fluctuation which renders the Maxwell term ineffective. Up to irrelevant couplings, we can safely set $a_\mu = A_\mu^{eff}$ in the above equation and find

$$\mathcal{L} = \psi_a^\dagger \gamma_\mu (\partial_\mu - i 2 \theta A_\mu) \psi_a + |(\partial_\mu - i 2 A_\mu) \phi|^2 + V(|\phi|) + \dots \quad (47)$$

where $\dots$ means the irrelevant couplings between bosons and fermions.

The above Eq. leads to in-consistent conclusion that the original charge neutral spinon ψ becomes charged: carrying charge e for $\theta = 1/2$ and $-e$ for $\theta = -1/2$. Therefore we also reach ambiguous conclusions for $\theta = 1/2$ and $\theta = -1/2$. The reason is that we replace the Z_2 CS gauge theory on the lattice which explicitly respects the equivalence between $\theta = 1/2$ and $\theta = -1/2$ by the $U(1)$ CS theory which does not respect this equivalence in the continuum limit. Identifying a_μ^ψ which is a discrete Z_2 gauge field with A_μ which is a $U(1)$ gauge field is a direct consequence of this problematic replacement.

B. Disordered and superconducting phases

We follow the discussions in the previous section. In the disordered phase, the vortex condense $\langle \Phi \rangle = \Phi_0$ which generates a mass term for a_μ^Φ in Eq.41. Integrating out the massive a_μ^Φ leads to a Maxwell term for a_μ^ψ .

$$\mathcal{L} = \psi_a^\dagger \gamma_\mu (\partial_\mu - i a_\mu^\psi) \psi_a + \frac{1}{4\Phi_0^2} (f_{\mu\nu}^\psi)^2 + \frac{1}{2} (a_\mu^\psi / \theta + A_\mu^{eff})^2 \quad (48)$$

The Dirac fermions become free and carry spin $1/2$ and charge $2\theta e$. In fact, we reach the same conclusion from the boson representation Eq.46.

In the superconductor phase, the vortex Φ is massive, therefore can be integrated out, it leads to the old Maxwell term $\frac{1}{4m_\Phi} (f_{\mu\nu}^\Phi)^2$. Integrating out a^Φ generates a mass term for a_μ^ψ :

$$\mathcal{L} = \psi_a^\dagger \gamma_\mu (\partial_\mu - i a_\mu^\psi) \psi_a + \frac{m_\Phi}{2} (a_\mu^\psi)^2 + \frac{1}{2} (a_\mu^\psi / \theta + A_\mu^{eff})^2 \quad (49)$$

where m_Φ is the mass of the vortex.

Diagonalizing the last two mass terms leads to a *continuously changing charge*. In fact, we reach the same description from the boson representation Eq.46. Because in superconductor phase, the boson condense $\langle \phi \rangle = \phi_0$, therefore generates a mass term for a_μ :

$$\mathcal{L} = \psi_a^\dagger \gamma_\mu (\partial_\mu - i \theta a_\mu) \psi_a + \frac{\phi_0^2}{2} (a_\mu)_t^2 + \frac{1}{2} (a_\mu + A_\mu^{eff})^2 \quad (50)$$

Which is essentially the same as Eq.49 after the identification $m_\Phi \sim \phi_0^2$.

Again, the pathological claims are due to the artifact of replacing the Z_2 mutual statistical gauge theory by a $U(1)$ mutual CS theory, then treating Z_2 gauge field and $U(1)$ electrical gauge field on the same footing. Although all these results are the artifacts of this replacement, they do suggest that the mutual statistical interaction between $hc/2e$ vortex and spinons leads to the confinement of spinon and chargon into electron and Cooper pair.

V. DISCUSSIONS AND CONCLUSIONS

The original singular gauge transformation [3,4] was proposed for static vortices. In this paper, we extend the singular gauge transformation to moving vortices generated by quantum fluctuation. By making a close analogy to the conventional singular gauge transformation of FQH system, we perform singular gauge transformations attaching flux of moving vortices to quasi-particles or vice versa. Just like conventional singular gauge transformation leads to conventional CS term, the two mutual singular gauge transformations lead to mutual CS term. In this way, we propose a intuitive and physical transparent approach to bring out explicitly the underlying physics associated with the condensation of the $hc/2e$ vortices.

Based on the earlier work by Balents, Fisher and Nayak [13], Senthil and Fisher developed Z_2 gauge theory [10,11] to study quasi-particles coupled to vortices generated by quantum fluctuations. By breaking electrons and Cooper pairs into smaller constituents: chargons with charge e ,

spin 0 and spinons with charge 0, spin 1/2, SF introduced a local Z_2 gauge degree of freedom to constrain the Hilbert space to the original one. The effective action describes both chargons and spinons coupled to local fluctuating Z_2 gauge theory with a doping dependent Berry phase term. By the combination of standard duality transformation of 3 dimensional XY model and that of Z_2 gauge theory, the action is mapped into a dual vortex representation where the $hc/2e$ vortices and spinons are coupled by a mutual Z_2 CS gauge theory. As usual vortices in XY model, the $hc/2e$ vortices also couple to a fluctuating $U(1)$ gauge field which mediates the long-range logarithmic interaction between the vortices. Starting from the dual representation, the authors in Ref. [11] studied a transition from d-wave superconductor to confined Mott insulator driven by the condensing of $hc/2e$ vortices. In order to study the critical behaviors of this particular confinement and de-confinement transition, they replaced the Z_2 mutual CS theory on the lattice by $U(1)$ mutual CS theory in the continuum and performed renormalization Group (RG) analysis. Unfortunately, some of their RG analysis are incorrect as demonstrated in Sec. IV.

It was shown in the text that our resulting general effective action Eq. 19 at the two special statistical angles $\theta = \pm 1/2$ is essentially equivalent to the dual vortex representation of Z_2 gauge theory developed by SF, except it also bring out explicitly the Volovik effect which was missed in SF's Z_2 gauge theory.

As explicitly stressed in this paper, the $U(1)$ mutual CS theory in the continuum does not have the required periodicity under $\theta \rightarrow \theta + 1$. In order to preserve this periodicity when the mutual statistical angle takes special values $\theta = \pm 1/2$, a Z_2 mutual CS theory must be enforced on the lattice. Again, two ways of regularization may lead to different conclusions. In fact, the periodicity of the conventional CS theory under $\theta \rightarrow \theta + 2$ is also a very intricate issue. On the one hand, the perturbative RG expansion in terms of the statistical angle θ in the continuum limit in Refs. [21–23] does not have this periodicity. On the other hand, properly regularized on the lattice, the CS theory does have this periodicity [28]. The two different regularization do lead to different conclusions on the Quantum Hall transitions. It also leads to notorious Hall conductivity difficulty at $\nu = 1/2$ [29]. Although we are unable to solve the critical behaviors of this combined Z_2 and $U(1)$ gauge fields coupled to spinons and $hc/2e$ vortices, we stressed the importance of putting the theory on the lattices to keep the periodicity of Z_2 mutual CS term and shed considerable lights on the physical picture and structure of the quantum transitions driven by $hc/2e$ vortices. We suggest that it may be possible to trace out the Z_2 gauge fluctuation on the lattice first to get an effective action without the discrete gauge fields, then take continuum limit of it and perform RG calculation. Further work is needed to sort out all the

possible phases and to understand the critical behaviors of the phase transitions between these phases.

This work was supported by NSF Grant No. PHY99-07949, university of Houston and DMR-97-07701. We thank M. P. A. Fisher, S. Girvin, S. Kivelson, P. A. Lee, S. Sachdev, T. Senthil, Z. Tسانovic, Y. S. Wu and Kun Yang for helpful discussions.

APPENDIX A: THE MOST GENERAL FORM OF THE GAUGE PROPAGATORS

Adding the gauge fixing terms $\frac{1}{2\alpha}((\partial_\mu a_\mu^\psi)^2 + (\partial_\mu a_\mu^\Phi)^2)$ to Eq. 22, we can find the gauge field propagators by inverting the matrix

$$\begin{pmatrix} \Pi_1(k)k(\delta_{\mu\nu} - \frac{k_\mu k_\nu}{k^2}) + \frac{k_\mu k_\nu}{\alpha} & \Pi_2(k)\epsilon_{\mu\nu\lambda}k_\lambda \\ \Pi_2(k)\epsilon_{\mu\nu\lambda}k_\lambda & \Pi_3(k)k(\delta_{\mu\nu} - \frac{k_\mu k_\nu}{k^2}) + \frac{k_\mu k_\nu}{\alpha} \end{pmatrix} \quad (A1)$$

The results are:

$$\begin{aligned} (G_{\psi\psi})_{\mu\nu} &= \frac{\alpha k_\mu k_\nu}{k^4} + \frac{\Pi_3}{\Pi_1\Pi_3 + \Pi_2^2} \frac{1}{k} (\delta_{\mu\nu} - \frac{k_\mu k_\nu}{k^2}) \\ (G_{\psi\Phi})_{\mu\nu} &= (G_{\Phi\psi})_{\mu\nu} = \frac{\Pi_2}{\Pi_1\Pi_3 + \Pi_2^2} (-\frac{\epsilon_{\mu\nu\lambda}k_\lambda}{k^2}) \\ (G_{\Phi\Phi})_{\mu\nu} &= \frac{\alpha k_\mu k_\nu}{k^4} + \frac{\Pi_1}{\Pi_1\Pi_3 + \Pi_2^2} \frac{1}{k} (\delta_{\mu\nu} - \frac{k_\mu k_\nu}{k^2}) \end{aligned} \quad (A2)$$

In the Landau gauge $\alpha = 0$ and putting $\vec{k} = 0$, we recover the results calculated in Sec. III.

APPENDIX B: APPLICATION TO STATIC DISORDERED VORTEX ARRAY

In Ref. [6], Ye pointed out that the long range Logarithmic interaction between vortices suppress the superfluid velocity fluctuation, but does not affect the internal gauge field fluctuation. He concluded that the quasi-particle scattering from the random gauge field dominate over that from the superfluid velocity (the Volovik effect). In this appendix, by using the formalism developed in the main text, we provide additional evidence for this conclusion.

For random pinned vortices, the static vortices do not feel the statistical gauge field a_μ^Φ in Eq. 19. This corresponds to a mass term for this gauge field. Adding the mass term to the equation and integrating out a_μ^Φ , it generates a Maxwell term for a_μ^ψ [30]:

$$\begin{aligned} \mathcal{L} &= \psi_a^\dagger \gamma_\mu (\partial_\mu - i a_\mu^\psi) \psi_a + \frac{1}{4} (f_{\mu\nu}^\psi)^2 \\ &+ |(\partial_\mu - i a_\mu) \Phi|^2 + V(|\Phi|) + \frac{1}{4} f_{\mu\nu}^2 - i A_\mu^{eff} \epsilon_{\mu\nu\lambda} \partial_\nu a_\lambda \end{aligned} \quad (B1)$$

In the above Eq. the averages over a_μ^ψ, a_μ, Φ should be understood as quenched instead of annealed averages. As discussed in the main text, the long-range logarithmic interaction between vortices represented by the Maxwell term suppress the fluctuation of the superfluid velocity. The quasi-particle is moving in a long-range correlated random magnetic field and short-range correlated scalar potential, in consistent with the physical picture in Ref. [6].

APPENDIX C: HC/E VORTICES, SPIN-CHARGE SEPARATION AND MOTT INSULATOR

In sec III, neglecting the charge fluctuation, we concentrated on the effect of mutual C-S term. In this appendix, we do the opposite, namely neglecting the mutual C-S term, but concentrate on the effect of charge fluctuation. This is realized by using neutral-like transformation, namely setting $\phi_A = \phi_B = \phi/2$ which is single-valued in the resulting hc/e vortices. The double strength vortex and its stability was investigated in Ref. [12], the associated spin-charge separation has been discussed extensively in Refs. [13,10]. In this appendix, to contrast with $hc/2e$ vortices discussed in the main text, we review their results, fill in some details and also get the new result on the expression of current operator I_μ in the superconducting phase.

Putting $\phi_A = \phi_B = \phi/2$, namely setting $a_\mu = 0$ in Eq.11 leads to:

$$\mathcal{L} = \psi_{1a}^\dagger [\partial_\tau + v_f p_x \tau^3 + v_2 p_y \tau^1] \psi_{1a} + (1 \rightarrow 2, x \rightarrow y) + \frac{K}{2} (\partial_\mu \phi - A_\mu^{eff})^2 \quad (C1)$$

where the effective gauge field is $A_\mu^{eff} = A_\mu - K^{-1} J_\mu$. There is no internal gauge field a_μ , because double vortices do not scatter the quasi-particles.

Performing duality transformation to vortex representation leads to

$$\mathcal{L} = \psi_a^\dagger \gamma_\mu \partial_\mu \psi_a + |(\partial_\mu - i a_\mu) \Phi|^2 + V(|\Phi|) + \frac{1}{4} f_{\mu\nu}^2 - i A_\mu^{eff} \epsilon_{\mu\nu\lambda} \partial_\nu a_\lambda - \mu \epsilon_{ij} \partial_i a_j \quad (C2)$$

where the last (Berry phase) term can be absorbed into A_μ^{eff} by redefining $A_\mu^{eff} \rightarrow A_\mu^{eff} + i\mu\delta_{\mu 0}$. Comparing with Eq.19, there is only charge fluctuation. There is no mutual C-S term, namely $\theta = 0$ or any *integer*.

The duality transformation to boson representation is

$$\mathcal{L} = \psi_a^\dagger \gamma_\mu \partial_\mu \psi_a + |(\partial_\mu - i a_\mu^\phi) \phi|^2 + V(|\phi|) + \frac{1}{4} (f_{\mu\nu}^\phi)^2 + i a_\mu \epsilon_{\mu\nu\lambda} \partial_\nu (a_\lambda^\phi - A_\lambda^{eff}/\theta) + \frac{1}{4} f_{\mu\nu}^2 \quad (C3)$$

Integrating out the electric gauge field leads to

$$\mathcal{L} = \psi_a^\dagger \gamma_\mu \partial_\mu \psi_a + |(\partial_\mu - i a_\mu^\phi) \phi|^2 + V(|\phi|) + \frac{1}{4} (f_{\mu\nu}^\phi)^2 + \frac{1}{2} (a_\mu^\phi - A_\mu^{eff})_t^2 \quad (C4)$$

Setting $a_\mu^\phi = A_\mu^{eff}$ up to irrelevant term leads to

$$\mathcal{L} = \psi_a^\dagger \gamma_\mu \partial_\mu \psi_a + n_0 \phi^\dagger \partial_\tau \phi + |(\partial_\mu - i A_\mu) \phi|^2 + V(|\phi|) + \dots \quad (C5)$$

where the second term is the Berry phase term and $\dots$ means the irrelevant couplings between bosons and fermions. The above Eq. is essentially the same as Eq.C1. Bosons and fermions are asymptotically decoupled. The fermions (spinons) carry only spin 1/2, the bosons (holons) carry only charge e . There is spin-charge separation. The long-range Coulomb interaction could be incorporated by adding the dynamic term for the time component of the gauge field $\frac{1}{2} k |A_0(k)|^2$ [31]. Without the fermionic part, the action is the same as the superconductor-insulator transitions in dirty metals studied in [32], except here condensed is a charge e boson instead of a Cooper pair.

In the underdoped regime, the vortex condenses $\langle \Phi \rangle = \Phi_0$ which generates a mass term for a_μ in Eq.C2 $\frac{1}{2} \Phi_0^2 (a_\mu)_t^2$. Integrating out the massive a_μ leads to a Maxwell term for $A_\mu^{eff} = A_\mu - K^{-1} J_\mu$ from which we can identify the automatically conserved electric current $I_\mu = \frac{1}{\Phi_0^2} (\partial_\nu^2 J_\mu - \partial_\mu \partial_\nu J_\nu)$ [13].

In the superconductor phase, the vortex Φ is massive, therefore can be integrated out, it leads to the old Maxwell term $\frac{1}{4m_\Phi} (f_{\mu\nu})^2$. Integrating out a_μ generates a mass term $\frac{1}{2} m_\Phi (A_\mu^{eff})_t^2$. From the mass term, we can identify the automatically conserved electric current $I_\mu = m_\Phi (J_\mu)_t$. Note that although the quasi-particle electric current is not conserved itself, the total electric current $I_\mu = m_\Phi (\delta_{\mu\nu} - \frac{q_\mu q_\nu}{q^2}) J_\nu(q)$ is conserved.

-
- [1] G. E. Volovik, Sov. Phys. JETP 58, 469 (1993)
 - [2] C. Kubert and P. J. Hirschfeld, Phys. Rev. Lett. 80, 4963 (1998).
 - [3] P. W. Anderson, cond-mat/9812063.
 - [4] M. Franz and Z. Tesanovic, Phys. Rev. Lett. 84, 554 (2000); O. Vafek *et al*, cond-mat/0007296.
 - [5] Luca Marinelli, B. I. Halperin and S. H. Simon, Phys. Rev. B 62, 3488 (2000).
 - [6] Jinwu Ye, Phys. Rev. Lett. 86, 316 (2001).
 - [7] K. Krishana *et al*, Science 277, 83 (1997), Phys. Rev. Lett. 82, 5108 (1999); H. Aubin *et al* Science 280, 9a (1998), Phys. Rev. Lett. 624, 5108 (1999).
 - [8] C. Dasgupta and B. I. Halperin, Phys. Rev. Lett. 47, 1556 (1981); X. G. Wen and A. Zee, Int. J. Mod. Phys. B 4,437 (1990).

- [9] M. P. A. Fisher and D. H. Lee, Phys. Rev. B39, 2756 (1989).
- [10] T. Senthil and M. P. A. Fisher, cond-mat/9910224, 9912380,0006481,0006500,0008082.
- [11] C. Lannert, M. P. A. Fisher, T. Senthil, cond-mat/0007002.
- [12] Jinwu Ye and S. Sachdev, Phys. Rev. B44, 10173 (1991); S. Sachdev, *ibid.* 45, 380 (1992); P. A. Lee and N. Nagaosa, *ibid.* 46, 5621 (1992).
- [13] L. Balents, M. P. A. Fisher and C. Nayak, Int. J. Mod. Phys. B12, 1033(1998), Phys. Rev. B60, 1654 (1999), *ibid* 61, 6307 (2000). For a review, see M. P. A. Fisher, cond-mat/9806164.
- [14] In this paper, we neglect the curvature term which may be important for the thermal Hall conductance in the presence of external magnetic field. For details, see Ref. [6].
- [15] We can also directly add Eq.2 and Eq.4 without performing this singular gauge transformation and then take continuum limit. But the resulting continuum action breaks spin $SU(2)$ symmetry.
- [16] In the impurity case, the inter-node scatterings maybe important because the quasi-particle momenta need not be conserved in the impurity scattering. see A. C. Durst and P. A. Lee, Phys. Rev. B 62, 1270 (2000). We do not consider the possible magnetic ordering transition connecting different nodes in this paper.
- [17] If the quasi-particle is hole-like, then $a_\mu \rightarrow -a_\mu$, but v_μ which is gauge invariant stays the same. We reach the same conclusion.
- [18] Note that under this transformation, the Time (T) reversal symmetry of c becomes Particle and Hole (PH) symmetry of d and vice versa. In the following, we use the language in the d representation.
- [19] B. I. Halperin, P. A. Lee and N. Read, Phys. Rev. B47, 7312 (1993).
- [20] W. Chen, M. P. A. Fisher and Y. S. Wu, Phys. Rev. B48, 13749 (1993).
- [21] X. G. Wen and Y. S. Wu, Phys. Rev. Lett. **70**, 1501 (1993).
- [22] G. W. Semenoff, P. Sodano and Y. S. Wu, Phys. Rev. Lett. **62**, 715 (1989); W. Chen, Phys. Lett. B251, 415 (1990).
- [23] Jinwu Ye and S. Sachdev, Phys. Rev. Lett, 80, 5409 (1998); Jinwu Ye, Phys. Rev. B60, 8290 (1999).
- [24] M. Campostrini *et al*, cond-mat/0010360.
- [25] T. Appelquist *et al*, Phys. Rev. Lett, 60, 2575 (1988); D. Nash, *ibid*, 62, 3024 (1989).
- [26] Yong Baek Kim and A. J. Millis, Physica, Vol. A4, 171 (1999); I. Ussishkin and A. Stern, Phys. Rev. B56, 4013 (1997); S. Sakhi, *ibid.* 56, 4098 (1997); Kun Yang, *ibid.* 58, R4246 (1998); Y. Oreg and B. I. Halperin, *ibid.* 60, 5679 (1999).
- [27] In double layer Quantum Hall systems, due to the T symmetry breaking by the external magnetic field, there is also a conventional CS term (the electron in a given layer sees its own flux). The K matrix [26] contains a zero eigenvalue, it indicates a Goldstone mode which has direct experimental consequences. See I. B. Spielman, *et al*, Phys. Rev. Lett, 84, 5808 (2000); cond-mat/0012094. In Eq.22, the K matrix only has off-diagonal elements, therefore invertible.
- [28] E. Fradkin and S. Kivelson, Nucl. Phys. B474, 543 (1996).
- [29] S. Kivelson *et al*, Phys. Rev. B55, 15552 (1997).
- [30] Strictly speaking, it is a *random* Z_2 gauge field.
- [31] M. P. A. Fisher and G. Grinstein, Phys. Rev. Lett, 60, 208 (1988); Jinwu Ye, Phys. Rev. B58, 9450 (1998).
- [32] M. C. Cha *et al*, Phys. Rev. B44, 6883 (1991); M. Wallin *et al*, *ibid* 49, 12115 (1994).